DOI: 10.30994/sjik.v10i1.797

ISSN: 2252-3847 (print); 2614-350X (online) Vol.10 No.1 May 2021 Page. 1348-1355

Analysis Risk Factors For Suicidal Thoughts Among Students In Surakarta: A Case Study Approach

Siti Khadijah¹, Sumardino¹, Ah Yusuf², Hanik Endang Nihayati²

¹Health Polytechnic of Surakarta, Ministry of Health Indonesia, Indonesia

²Universitas Airlangga Surabaya, Indonesia

* Correspondent Author: s.khadijah3029@gmail.com

ABSTRACT

The design of this study was a qualitative case study with eight respondents who were selected using the purposive sampling technique. The inclusion criteria were students with a positive risk of suicide and high scores at the screening phase. Data were obtained by semi-structured interviews and analyzed thematically using the Colaizzi method. This study showed that the exploration of the risk factors for suicidal thoughts consisted of two themes, eight sub-themes, 25 categories, and 50 keywords. Two themes are suicide, and spiritual. Eight sub-themes namely Causes, Response to problems, Family attitude, Efforts that have been made, Obstacle factors, Supporting factors, Meaning of the problem faced, A feeling that arose spiritually. The risk factors for suicidal thoughts among students in Surakarta are that they have no interest in entering higher education, have difficulty in attending lectures, disappointing academic scores, broken home families, problems with a partner, financial issues, sadness, and disappointment in God, and they face problems repeatedly.

Keywords: Risk Factors, Suicidal Thought, College Students

Received March 17, 2021; Revised April 7, 2021; Accepted April 28, 2021



TRADA Jurnal Ilmiah Kesehatan, its website, and the articles published there in are licensed under a Creative Commons Attribution-ShareAlike 4.0 International License.

Website: https://sjik.org/index.php/sjik | Email: publikasistrada@gmail.com 1348

DOI: 10.30994/sjik.v10i1.797

ISSN: 2252-3847 (print); 2614-350X (online) Vol.10 No.1 May 2021 Page. 1348-1355

BACKGROUND

Students should have a fighting mentality and tremendous enthusiasm as agents of social change to make improvements for the better while avoiding the detrimental consequences of modernization (Bestari, 2015). At this time, the highest prevalence of suicide is among young adults, including students. Suicide is the second global major cause of death for people aged 15-29 years (young adults), and most students are in this age range (WHO, 2016). The young adulthood process is the phase of the transition of a person from puberty to adulthood. Someone who typically focuses on himself, becomes someone who can think and prepare for the future. This young adulthood process varies from facing external and internal challenges such as mental health issues (Viora, 2018). This adulthood phase is also considered as an adaptation to life and a new social expectation. Being adaptable to a social expectation means developing and changing new attitudes, desires, and values to fulfill their roles. Difficulty to adapt in this young adult phase is somewhat challenging (Pieter and Lubis, 2017). Suicide risk is the risk of attempting to harm oneself to end life (PPNI, 2016).

Based on the phenomenon in the Higher Education Department of Nursing and Midwifery in Ghana, a primary study said there were reports of suicide among college students (Nii *et al.*, 2019). The study of (Wolitzky-taylor *et al.*, 2019) revealed that worldwide suicides in 2018 resulted in 800,000 deaths. World Health Organization (WHO) in 2016 showed that the suicide rate in Indonesia continues to increase, from 4.3% in 2012 to 5.2% in 2016. In 2019, a student at a university in Surakarta committed suicide (Ryantono, 2019), and another one in Bandung (Anonim, 2019). Apart from these two cases, there were still many suicides that occurred among students. Nationally, Central Java is the province with the highest number of suicides with 331 cases, followed by East Java with 119 cases. A preliminary study on students at the Health Polytechnic of Surakarta in 2019 showed 32% of individuals in high suicide ideas and 68% in low suicide ideas.

Based on the preliminary study, which showed that the idea of suicide among students is still high, it should be well-considered because it will affect the students' fighting mentality. It is important to keep them working and studying productively, and early detection of risk factors of suicidal thoughts in students should be initiated. Hence, it is necessary to explore the overview of the risk factors for suicidal thoughts among students so that suicide prevention efforts can be conducted accordingly. Aim; The purpose of this study was to discover the risk factors for suicidal thoughts and desires among students in Surakarta, Indonesia.

METHODS

This study used a qualitative case study design. The study participants consisted of first-year students at the Health Polytechnic of Surakarta who were young adults aged 17-21. Eight participants were selected using a purposive sampling technique with a positive risk of suicide and high scores at the screening phase. Interviews were conducted from December 2019 to January 2020 with semi-structured interviews. The data collection tool was a recording device, and data were analyzed thematically using the Colaizzi method. Ethical feasibility for this research was carried out at the ethics committee at the Faculty of Nursing, Airlangga University on December 9, 2019, Ethics certificate number 1854-KEP.

DOI: 10.30994/sjik.v10i1.797

ISSN: 2252-3847 (print); 2614-350X (online) Vol.10 No.1 May 2021 Page. 1348-1355

RESULTS

Based on the results of semi-structured interviews with eight students, it was found that both internal and external factors caused a suicidal thought or idea. This study was a qualitative case study design with the characteristics of the respondents as follows:

Table 1 Characteristics of respondents in phase one

No	Code	Gender	Age	Level of education
1	M1	Male	17	Diploma III
2	M2	Female	18	Diploma IV
3	M3	Female	20	Diploma IV
4	M4	Female	20	Diploma III
5	M5	Female	20	Diploma IV
6	M6	Female	21	Diploma III
7	M7	Female	19	Diploma III
8	M8	Female	18	Diploma III

Table 1 describes the characteristics of the respondents. There were male and female first-year students with an age range of 17-21 years old involved in the study. A purposive sampling technique was used to select the participants with a positive risk of suicide and high scores at the screening phase. The majority of respondents were female.

Based on the analysis, there were two themes, eight sub-themes, 25 categories, and 50 keywords. The themes were identified from sub-themes, the sub-themes were concluded from the categories, and the categories were summarized from the keywords. The overall themes, sub-themes, categories, and keywords are as follows:

Theme 1: Suicide with a sub-theme of 1 cause; category 1 does not match specialization, keywords; I don't like being forced, not according to my passion. Category 2 damaged houses, keywords; disappointed, sad, parents are not in the same house, one parent is cheating, the family is not harmonious often quarrels, one house with the stepfather and the relationship is not good. Category 3 lover, keywords; abandoned by a lover, an authoritarian lover, afraid of losing. Category 4 economy, keywords; has to help a lot, withstand the lack of conditions, limited facilities. Category 5 academic, the keyword is difficulty in understanding the lesson, the value is not as expected. Category 6 Bullying, keywords; embarrassed to be embarrassed in front of the crowd. Sub-theme 2 Changes in behavior/response to problems; 7 moody categories, keywords; prefers to be alone, away from parents, to be more introverted. Category 8 speak as needed, keywords; in trust, disappointed with the parents. Category 9 Crying, keywords; sad about related problems. Category 10 is not passionate in any way, keywords; lazy to go to college, have no appetite, have trouble sleeping or sleep a lot. Sub-theme 3 Efforts made; category 11 confide in friends, keywords fear of being scolded by parents, feeling grown-up, more comfortable. Category 12 confide in family, keywords; to convey it to parents. Category 13 confide in spiritual mentors, keywords; got strength spiritually. Sub-theme 4 family attitudes; category 14 do not care, keywords are often angry, insinuating, comparing with relatives or other people. Sub-theme 5; obstacle factor; 15 categories, friends, keywords; insult, Category 16 family, keywords; less harmonious. Category 16 Thoughts, keywords; negative thoughts arise. Sub-theme 6 Supporting factors; category 17 motivation, keywords; self-motivation, family motivation, lover motivation, friend motivation.

Theme 2; spiritual with sub-theme 1 feeling that arises spiritually; category 1 disappointed, keywords; let go, be grateful. Category 2 sad, keywords; patient. Category 3 angry, keywords; forgive, start loving yourself, don't judge yourself. Sub-theme 2 the

DOI: 10.30994/sjik.v10i1.797

ISSN: 2252-3847 (print); 2614-350X (online) Vol.10 No.1 May 2021 Page. 1348-1355

meaning of the problem, category 4 have considered as punishment, keywords; there have been mistakes in the past. Category 5 feels God is unfair, keywords; was given a problem in a row. Category 6 considers it a disaster, keywords; in life, there must be trials from God.

Table 2 Risk factors for suicidal thoughts among students

Theme 1: Suicide

Subtheme	Significant Statements
Causes	"I went to college not according to my interest, so it felt difficult to attend lectures" (P1, P2, P4, P5, P7, P8). "I have suicidal thoughts because I have
	a broken home family, Parents who are not in a harmonious relationship
	(P2, P4, P5, P6, P7), "financial issues become a heavy burden and must
	be forced to many limitations so that it weighs on my mind(P2, P7, P8),
	"withstand many limitations" (P7, P8), "Suicidal thoughts arise because of an authoritarian partner and a partner's abandonment" (P4, P6),
	" I cannot stand with the trauma of being bullied by friends" (P3, P8),"
	grades that do not match expectations, difficulty taking lessons and having
	low academic scores, and feeling depressed because of shame" (P5, P8).
	Each student has more than one problem simultaneously and repeatedly,
	resulting in frustration and triggering suicidal thoughts.
Response to	"when disappointed with parents, I am lazy to talk to parents and family
problems	members, I become more moody, introverted, and not enthusiastic (P1, P2, P2, P5, P6, P7, P8)
	P3, P5, P6, P7, P8), "problems make me overthinking, lack of appetite, and having trouble with sleeping." (P3, P4, P6). " being look of self-genfidence."
	having trouble with sleeping" (P3, P4, P6), "being lack of self-confidence and trust to others" (P3, P6), "with the problem I have, I feel stuck so I
	prefer to be alone in the room" (P1).
Efforts that	"the efforts made were playing to a friend's house and telling them "(P1,
have been	P2, P4, P5, P7), and "told my aunt and told my parents but still felt that they
made	did not provide a solution to continue studying even if it was being forced"
	(P8)," the problem buried itself" (P6, P3).
Family	"families often get angry" (P1, P5, P7, P8), " the family does not care
attitude	about the problems experienced by its children" (P2, P6, P8), " families often compare with other siblings" (P1), " the family often brings up and
	insinuates about the problems that exist in the family" (P5).
Obstacle	"friends who sometimes bully" (P1, P8), "family conditions that are not
factors	harmonious, parents who are often angry and often disputing" (P2, P5, P8),
	"always being overthinking" (P3, P4, P6, P7), "mistrust with friends"
	(P3, P6, P8).
Supporting	"personal motivation" (P1, P6), "motivation from siblings and
factors	cousins" (P2, P8), "motivation from partner" (P4, P5).

Theme 2: Spirituality

Subtheme	Significant Statements
Meaning of the	"assuming this is a punishment from God" (P1, P2, P4, P5), "this
problem faced	problem is part of a trial for me" (P3, P7), "think that God is unfair" (P6, P8).

DOI: 10.30994/sjik.v10i1.797

ISSN: 2252-3847 (print); 2614-350X (online) Vol.10 No.1 May 2021 Page. 1348-1355

A feeling that	"feelings of disappointment in God because I have been praying
arose spiritually	diligently but am still given repeated problems" (P1, P3, P4, P5),
	"feelings of sadness why am I burdened like this" (P2, P6, P7, P8)

DISCUSSION

Identifying the risk factors for suicidal thoughts as early as possible could minimize the incidence of suicide risk. In the theme of suicide, the risk factors for suicidal thoughts among students found were they enter the college with a major that is not what they want, broken home or issues in the family, loss of a partner, and academic problems. If the individual does not have a good coping mechanism or is adaptive, it will make a person experience psychological pressure and impact the risk of suicide. Factors that influence suicide according to Yosep, (2011), one of which is social isolation and human relations. Suicide, in general, is due to the stress that arises because of failure to be adaptive. This situation could happen in the work environment, family, school, society relationship, etc. Likewise, when someone feels isolated, losing a relationship, or being cut off from loved ones. Yusuf, Fitriyasari, R, and Nihayati (2015), clarified that suicide is caused by many causes, one of which is psychosocial factors, such as loss, lack of social support, negative life events, chronic physical illness, and a family history of suicide.

Park (2017), explained that the family function has a significant correlation with mental health, depression, and suicidal ideation on freshmen students at two universities. Related research, Andari (2017) discovered that the suicide case in Gunung Kidul is mostly the result of depression and chronic illness. Depression can occur due to loss of jobs, not affording family needs, divorce, infidelity and breakups, and chronic diseases are common causes of suicide in older people. Mehl-madrona (2016) described that the incidence of suicide is based on three themes: 1) sudden, unwanted presence of a third person; 2) public humiliation of others followed by a high degree of shame; and 3) stress levels due to poverty.

Furthermore, Tumon (2014), explained that bullying has a tendency towards depression in adolescents, where the depression results in thoughts of committing suicide and self-injury. Research by Neupane *et al.*, (2020) reported that bullying cases in Nepalese school adolescents resulted in mental health problems such as high loneliness, anxiety, attempted suicide, school absence because of fear. Lower socioeconomic status at the individual and regional level also has been shown to raise the likelihood of suicidal thoughts. The poverty rate is closely related to the suicide rate in both males and females, and alcohol involvement is also positively associated with suicide. The economic effect plays a significant role in increasing alcohol-related suicide (Pirkis *et al.*, 2017; Kerr *et al.*, 2017). Another study, Kartika (2018), found a significant relationship between academic stress and suicidal ideation. Research Lew *et al.*, (2020) explained that higher scores in psychological stress, depression, anxiety, and stress are psychological risk factors for suicide among undergraduate students in China. Demographic factors associated with a higher risk of suicide are female, younger age, poor academic results, an only child, not participating in school associations, and owning a home in urban areas.

The risk factors for suicidal thought among students that were related to the spiritual theme identified interpreting problems faced negatively. This significantly triggers suicidal thoughts because strong beliefs will make som Spiritual is something that is related to spirit, enthusiasm for belief, hope and meaning in life eone interpret a problem in life positively. As a result, it will prevent the existence of suicidal thoughts. Spiritual is something that is related to spirit, enthusiasm for belief, hope and meaning in life (Yusuf, Nihayati and Iswari, 2017). The study (Hirono, 2019) reported that explaining faith in a higher power gives life

DOI: 10.30994/sjik.v10i1.797

ISSN: 2252-3847 (print); 2614-350X (online) Vol.10 No.1 May 2021 Page. 1348-1355

purpose and meaning, provides comfort, and prevents suicide. Another research (Umma, 2017) defined a significant influence from the life events (stressors), opinions on suicidal treatment, social support, friendship, and religiosity to suicide ideation. Garroutte & Goldberg's research in (Shinde and Wagani, 2019) found that commitment to spirituality was significantly associated with decreased suicide attempts. Spirituality had the effect of converting negative thoughts into positive ones. Other related research by Walker, Shin, and Kaslow (2018) explained that spiritual well-being leads to a protective function against feelings of isolation that can reduce the risk of an individual's psychological negativity. As a result, it can increase spiritual sense and social connectivity (interpersonal relationships) and reduces suicidal ideation, hopelessness, and symptoms of depression. The risk factor for suicidal thoughts is not a single factor, but several factors cause suicide, and each factor is interrelated. Related research from (Blasco, Castellví, and Almenara, 2016) showed that it is vital to explore and identify factors that affect suicide risk to inform and improve further intervention strategies. Hence, interventions are more effective when they focus on risk factors and protective measures.

CONCLUSION

Students have a risk of suicide so that the risk factors for suicidal thoughts were identified among students, and two themes were found, namely suicide and spirituality. The risk factors for suicidal thoughts among students in Surakarta are that they have no interest in entering higher education, have difficulty attending lectures, disappointing academic scores, broken home families, problems with partner, financial issues, sadness and disappointment in God, and they face problems repeatedly.

NURSING IMPLICATION

Universities or nursing schools should prepare strategies in dealing with existing conditions to prevent suicide among students. Preventive efforts within the Educational Institution in the form of creating an environment that is conducive to mental health development, providing communication, education and providing psychosocial support by means of infrastructure, including sports facilities, worship facilities, counseling facilities so as to minimize psychosocial problems (Yusuf, Fitriyasari and Nihaayati, 2019).

ACKNOWLEDGMENT

The author would like to express his deepest gratitude to the Administration of Airlangga University in Surabaya, Poltekkes Kemenkes Surakata, and the Ministry of Health scholarships for the financial support provided in carriying out studies.

REFERENCES

Andari, S. (2017) 'The suicide phenomenon in the Gunung Kidul', 7(1), pp. 92–107.

Anonim (2019) 'Detik-detik Mahasiswa S2 ITB Gantung Diri Bikin Merinding, Maaf Aku Tidak Sanggup Lagi"", *TribunJabar*, September.

Bestari, P. (2015) Mahasiswa, 25 kiat mahasiswa menjawab tantangan era otonomi daerah, masyarakat ekonomi ASEAN hingga generasi Indonesia Emas 2045. Bandung: Alfabeta.

Blasco, M. J., Castellví, P. and Almenara, J. (2016) 'Predictive models for suicidal thoughts and behaviors among Spanish University students: rationale and methods of the UNIVERSAL (University & mental health) project', *BMC Psychiatry*. BMC Psychiatry, 16, pp. 1–13. doi: 10.1186/s12888-016-0820-y.

DOI: 10.30994/sjik.v10i1.797

ISSN: 2252-3847 (print); 2614-350X (online) Vol.10 No.1 May 2021 Page. 1348-1355

- Hirono, T. (2019) 'Preventing Soldiers and Veterans Suicide by Pastoral Counseling and Mental Health Treatment', *Journal of Pastoral Care & Counseling*, 73(3), pp. 176–182. doi: 10.1177/1542305019872433.
- Kartika, A. (2018) *Hubungan antara stres akademis dan ide bunuh diri pada mahasiswa*. Universitas Sanata Dharma Yogyakarta.
- Kerr, W. C. *et al.* (2017) 'Economic Recession, Alcohol, and Suicide Rates_ Comparative Effects of Poverty, Foreclosure, and Job Loss', *American Journal of Preventive Medicine*. Elsevier Inc., 52(4), pp. 469–475. doi: 10.1016/j.amepre.2016.09.021.
- Lew, B. *et al.* (2020) 'Suicidality among Chinese college students: A cross-sectional study across seven provinces', 15(8), pp. 1–13. doi: 10.1371/journal.pone.0237329.
- Mehl-madrona, L. (2016) 'Indigenous Knowledge Approach to Successful Psychotherapies with Aboriginal Suicide Attempters', *The Canadian Journal of Psychiatry*, 61(11), pp. 696–699. doi: 10.1177/0706743716659247.
- Neupane, T. *et al.* (2020) 'Correlates of bullying victimization among school adolescents in Nepal: Findings from 2015 Global School-Based Student Health', pp. 1–13. doi: 10.1371/journal.pone.0237406.
- Nii, E. *et al.* (2019) 'Prevalence of suicidal behaviour among nursing and midwifery college students in Ghana', (July 2018), pp. 1–10. doi: 10.1002/nop2.271.
- Park, Y. (2017) 'Influence of Family Function, Mental Health and Depression on Suicidal Ideation in Fresh College Joiners', 20, pp. 5065–5079.
- Pieter, H. and Lubis, N. . (2017) *Pengantar Psikologi dalam Keperawatan*. 3rd edn. jakarta: PT Kharisma Putra Utama.
- Pirkis, J. et al. (2017) 'Socio-Economic Position and Suicidal Ideation in Men', International Journal of Environmental Research and Public Health, 14(365). doi: 10.3390/ijerph14040365.
- PPNI (2016) Standar Diagnosis Keperawatan Indonesia. 1st edn. jakarta: PPNI.
- Ryantono (2019) 'Mahasiswa S2 UNS Solo Tewas Bunuh Diri, Selama Lima Bulan Terakhir Konsumsi Obat Depresi', *Tribun Solo*.
- Shinde, F. and Wagani, R. (2019) 'Does Spirituality Work as a Buffer in Suicide: A Systematic', 14(2), pp. 249–256.
- Tumon, M. (2014) 'Studi Deskriptif Perilaku', 3(1), pp. 1-17.
- Umma, A. (2017) Faktor Risiko Yang Memengaruhi Gagasan Bunuh Diri Pada Mahasiswa Kedokteran Preklinik Uin Syarif Hidayatullah Jakarta 2016-2017.
- Viora, E. (2018) 'Kesehatan Jiwa Remaja. Makalah disajikan dalam Temu Media 2018 HKJS'.
- Walker, A. L. G. K. L., Shin, L. J. and Kaslow, N. J. (2018) 'Spiritual well-being and psychological adjustment: mediated by interpersonal needs?', *Journal of Religion and Health*. Springer US, 57(4), pp. 1376–1391. doi: 10.1007/s10943-016-0275-y.
- WHO (2016) *Data Bunuh Diri*. doi: http://www.who.int/mental_health/ pencegahan / bunuh diri / suicideprevent / id /.
- Wolitzky-taylor, K. *et al.* (2019) 'Suicide prevention on college campuses: What works and what are the existing gaps? A systematic review and meta-analysis', *Journal of American College Health*. Taylor & Francis, 0(0), pp. 1–11. doi: 10.1080/07448481.2019.1577861.
- Yosep, I. (2011) Keperawatan Jiwa (edisi Revisi). Bandung: Refika-aditama.
- Yusuf, A., Fitriyasari, R, & and Nihayati (2015) *Buku Ajar Keperawatan Kesehatan Jiwa*. jakarta: Salemba Medika.

DOI: 10.30994/sjik.v10i1.797

ISSN: 2252-3847 (print); 2614-350X (online) Vol.10 No.1 May 2021 Page. 1348-1355

Yusuf, A., Fitriyasari, R. and Nihaayati, H. (2019) Kesehatan Jiwa Pendekatan Holistik Dalam Asuhan Keperawatan. Pertama. jakarta: Mitra Wacan Medika.

Yusuf, Nihayati and Iswari, M. . (2017) *Kebutuhan Spiritual: Konsep dan Aplikasi dalam Asuhan Keperawatan*. Edisi Pert. Jakarta: Mitra Wacan Media.